International Journal of Advanced Social Sciences and Education (IJASSE)
Vol. 3, No. 6, 2025: 425-436

# Qur'anic Embryology and Modern Developmental Physiology: A Comparative Analysis of Nutfah, 'Alaqah, Mudghah Stages with Contemporary Prenatal Biomarkers and Community Health Implications

HM. Hamdani[1]*, Subhan Muhith[2], Iman Tarmizi Taher[3], Rahma Zakiyah Al Hamdani[4], Melawati[5]
[1, 5]Universitas Aisyah Pringsewu
[2]Universitas Muhammadiyah Dahlan Palembang
[3]Universitas Sriwijaya Palembang
[4]Universitas Muhammadiyah Palembang
**Corresponding Author:** HM. Hamdani kangdani135@gmail.com

A R T I C L E I N F O

*Keywords: Qur'anic Embryology, Nutfah, 'Alaqah, Mudghah, Prenatal Biomarkers, Islamic Physiology, Maternal Health*

*Received: 19, October*
*Revised: 20, November*
*Accepted: 30, December*



A B S T R A C T

This study presents a comprehensive comparative analysis between Qur'anic descriptions of early human development specifically the sequential stages of nutfah, 'alaqah, and mudghah and contemporary biomedical knowledge on embryogenesis and maternal physiological biomarkers. Although the Qur'an refers to these stages in a concise linguistic form, classical and modern exegesis suggest that the terminology indicates distinct developmental phases. Modern embryology, on the other hand, describes the same period (weeks 1-8) using molecular, anatomical, and biochemical parameters such as fertilization, implantation, gastrulation, and early organogenesis. Using a PRISMA-guided systematic review combined with an integrative physiological analysis, this study synthesizes evidence from Qur'anic exegesis (classical and contemporary), peer-reviewed embryology articles, WHO developmental timelines, and hormonal biomarker studies (β-hCG, PAPP-A, progesterone, estrogen, AFP). A total of 68 eligible studies were analyzed after screening 1,242 titles. The study proposes a new framework, the Qur'anic Biophysiological Mapping Model, which integrates religious terminology and biomedical markers into a unified timeline. Findings hold significant implications for maternal health education in Muslim communities, especially in culturally sensitive counselling, prenatal awareness programs, and public-health communication. This interdisciplinary synthesis shows that faith-based concepts can harmoniously coexist with scientific explanations, offering a holistic view of early human development.

DOI: https://doi.org/10.59890/ijasse.v3i6.248
ISSN-E: 2988-2133
https://dmimultitechpublisher.my.id/index.php/ijasse

## INTRODUCTION

The study of human embryogenesis has historically been grounded in anatomical observation, physiological experimentation, and, in recent decades, molecular biology. However, the discourse on human development has long transcended scientific domains and entered the realm of religion, theology, and philosophy. One of the most compelling intersections emerges between Islamic scripture particularly the Qur'an and modern developmental physiology. Verses such as QS Al-Mu'minun 23:12-14 outline sequential stages of human creation using the terms nutfah, 'alaqah, and mudghah. These verses have been the subject of extensive exegetical, philosophical, and scientific analysis.

Despite the breadth of discussion, the literature exhibits notable fragmentation. Islamic studies typically focus on linguistic and theological interpretation, whereas biomedical sciences emphasize anatomical chronology and physiological mechanisms. Only a few modern works attempt to bridge the two domains meaningfully, and even fewer examine the convergence using measurable physiological parameters such as maternal biomarkers. Biomarkers like β-hCG, PAPP-A, progesterone, estrogen, and α-fetoprotein are crucial indicators of embryonic development and maternal adaptation, yet their relationship to Qur'anic staging has not been thoroughly investigated.

The need for interdisciplinary integration becomes more pressing within contemporary Muslim-majority societies, where prenatal health education often relies heavily on faith-based terminology. Bidan (midwives), nurses, and Islamic health educators commonly refer to nutfah, 'alaqah, and mudghah during counselling sessions. However, without a clear biomedical explanation that aligns with these terms, health promotion efforts can lack scientific grounding, reducing their effectiveness in improving maternal health literacy.

Furthermore, several global health issues underscore the importance of integrating culturally rooted concepts with biomedical knowledge. These include high rates of undetected early pregnancy complications, low awareness of hormonal changes in early gestation, and misunderstandings regarding the timing of embryonic development. Integrating Qur'anic terminology with developmental physiology could therefore enhance health communication, reduce misconceptions, and promote evidence-based maternal behaviors within Muslim communities.

From a scholarly standpoint, significant research gaps persist. Most previous publications merely compare the meanings of nutfah, 'alaqah, and mudghah descriptively, without connecting them to quantifiable physiological data. Virtually no study systematically maps Qur'anic staging with modern embryological weeks using standardized research frameworks such as PRISMA. Moreover, no existing research establishes correlations between Qur'anic stages and maternal biomarkers measured in clinical practice. This represents a major gap, as biomarkers serve as direct physiological indicators of developmental milestones.

## LITERATURE REVIEW

This study aims to fill these gaps with a structured, methodological, and evidence-based comparative analysis. By employing PRISMA for systematic literature selection, evaluating textual sources using established quality-assessment tools (CASP for religious/qualitative texts, NOS for biomedical studies), and synthesizing data through thematic and chronological mapping, this research provides a comprehensive interdisciplinary model. The resulting Qur'anic–Biophysiological Mapping Model offers a novel framework for understanding early human development through both religious and biomedical lenses. It is expected to benefit Islamic scholars, medical educators, clinicians, health promoters, and the broader academic community.

Thus, this paper seeks to answer three key research questions:

1. What are the linguistic, exegetical, and conceptual meanings of nutfah, 'alaqah, and mudghah in the Qur'an?
2. How do these stages correspond to modern embryology and the physiological changes experienced by pregnant women?
3. How can maternal biomarkers be integrated into a unified explanatory model aligned with Qur'anic terminology?

## METHODOLOGY

### Qur'anic Concept of Human Embryogenesis

The Qur'anic narrative of early human development is presented in several verses, most explicitly in Surah Al-Mu'minun (23:12-14), Al-Hajj (22:5), As-Sajdah (32:7-9), and Az-Zumar (39:6). These verses outline a sequential, phase-based development from nutfah to 'alaqah to mudghah, followed by bone formation ('idhām), flesh covering (lahm), and finally the transition into a distinct human being (khalqan ākhara). Unlike classical embryology texts, which describe development through anatomical or molecular milestones, the Qur'anic framework employs descriptive, metaphor-rich terminology that integrates biological, philosophical, and spiritual dimensions. This section explores these terms through linguistic analysis, classical tafsir, and contemporary scholarly interpretation.

### Nutfah

The term nutfah appears more than a dozen times in the Qur'an, referring literally to "a drop," "a small quantity of fluid," or "a mixed fluid" (nutfah amsyāj, QS 76:2). Classical scholars such as Ibn Kathir and Al-Tabari describe nutfah as a minute biological entity originating from both parents. Modern exegetes-including Sayyid Qutb, Al-Maraghi, and Shalih Al-Munajjid-interpret nutfah as the initial stage of conception, corresponding to fertilization and the earliest cellular divisions.

Biomedically, nutfah aligns with the period encompassing sperm, oocyte fusion, formation of the zygote, and cleavage divisions leading to the morula. This stage spans approximately days 1-5 post-fertilization. The description in Qur'an 76:2 of a "mixed drop" parallels the fusion of paternal and maternal gametes and the unique genomic recombination that occurs at syngamy. Further, Qur'an 53:46 ("Created from a fluid emitted") reflects the unidirectional contribution of sperm, while verses like Qur'an 4:1 ("Created from a single soul and its mate") allude to genetic complementarity and shared lineage.

**'Alaqah**

The term 'alaqah appears in the earliest revealed verses (QS 96:1-2), linking the beginning of human creation with this stage. Linguistically, 'alaq has three primary meanings:

1. Something that clings or attaches.
2. A suspended or hanging object.
3. A blood-like or clot-like substance. All three meanings correspond remarkably well with modern descriptions of implantation, trophoblastic invasion, and the early establishment of maternal embryonic circulation.

Classical tafsir including works of Al-Razi, Al-Qurtubi, and Ibn Ashur emphasize the clinging and dependent nature of the 'alaqah, drawing parallels with an organism that adheres to the uterine wall and draws nourishment from maternal blood. Modern embryology confirms that around days 6-13, the blastocyst attaches to and embeds itself within the endometrium, forming structures that "cling" tightly to maternal blood vessels. The formation of chorionic villi, lacunar networks, and early hematopoietic islands reflects the "blood like" description inherent in 'alaqah.

Moreover, the physiological spike of β-hCG during days 8-14, representing trophoblast activity, coincides temporally with the Qur'anic 'alaqah phase. This constitutes one of the key convergence points analyzed in this paper's mapping model.

**Mudghah**

The term mudghah literally means "a small chewed-like lump." Classical exegetes highlight the morphological resemblance of this stage to a piece of flesh bearing tooth mark like indentations. Modern embryology identifies this as the somite-formation stage approximately weeks 3-4, when segmented structures appear along the neural tube. These somites, arranged in repetitive blocks, confer a "chewed-like" appearance, making the Qur'anic description anatomically coherent.

From a physiological perspective, the mudghah stage marks the onset of primary organogenesis, including early cardiac looping, neural crest cell migration, and initial limb-bud formation. These developments align with the Qur'anic transition to subsequent phases: bone ('idhām) formation and muscular covering (lahm). Verses in Surah Al-Mu'minun (23:14) explicitly mention these phases sequentially, consistent with the biomedical timeline of osteogenesis followed by myogenesis.

**Integration of Qur'anic Descriptions with Developmental Phases**

The Qur'anic embryological framework, though concise, presents a clear chronological pattern. The sequencing of nutfah → 'alaqah → mudghah maps onto three major biomedical developmental clusters:

| No | Qur'anic Term | Approx. Biomedical Phase | Physiological Events |
|---|---|---|---|
| 1 | **Nutfah** | Fertilization → Cleavage (Days 1–5) | Zygote formation, genomic recombination, and early mitosis |
| 2 | **'Alaqah** | Implantation (Days 6– | Trophoblast invasion, lacunar |

| | | | |
|---|---|---|---|
| | | 14) | circulation, β-hCG surge |
| 3 | **Mudghah** | Early Organogenesis (Weeks 3–4) | Somite formation, neural tube closure, primitive heart |

The Qur'an's description of bones and muscles forming subsequently is also consistent with the sequence of embryonic tissue differentiation found in modern developmental biology.

**The 30 Month Gestational, Weaning Formula**

A unique Qur'anic contribution to developmental physiology is found in QS 46:15:

"His mother carried him and weaned him in thirty months."

Classical jurists (Ibn Abbas, Imam Malik, Ibn Qudamah) deduce from this verse and the 24-month breastfeeding mandate in QS 2:233 that the minimum viable gestational period is six months. This interpretation historically guided Islamic legal rulings on lineage and infant viability.

Biomedically, preterm infants around 24–28 weeks can survive with intensive care, aligning loosely with the classical six-month minimum when converted from lunar to solar months. While not identical, the proximity is notable, indicating that Qur'anic conceptual boundaries approximate modern viability thresholds.

**Conceptual Harmony Between Revelation and Physiology**

Rather than offering a technical embryology textbook, the Qur'an presents developmental stages that describe the essence, function, and relational nature of early human existence. Each stage emphasizes dependence, transformation, and divine intentionality. Modern physiology, although grounded in empirical observation, arrives at descriptions that match these conceptual stages. The overlap is significant enough to allow a comparative, integrative analysis without conflating scientific and theological domains.

## RESULT AND DISCUSSION

**Theme 1: Physiology of Early Pregnancy**

The physiology of early pregnancy represents a complex interplay between endocrine signaling, cellular differentiation, and maternal systemic adaptation. Early gestation consists of three pivotal events fertilization, implantation, and early organogenesis each corresponding closely with the Qur'anic phases nutfah, 'alaqah, and mudghah. These biological events are regulated by hormone surges, genomic activation, and embryonic–maternal communication that ensure viability and sustained development.

**Fertilization and Cleavage (Nutfah Phase)**

Within the first 24 hours after coitus, capacitated spermatozoa undergo the acrosome reaction, enabling penetration of the zona pellucida and fusion with the oocyte. This yields a diploid zygote the clearest biomedical analogue of nutfah. The zygote initiates synchronous mitotic divisions (cleavage) into blastomeres, forming a morula by day 3-4.

Genomic activation occurs around the 4-8 cell stage, aligning with Qur'anic descriptions of nutfah amsyāj ("mixed drop"), referring not only to parental fluid but also to genetic mixing. During this time, progesterone from the corpus luteum stabilizes the endometrium, while estrogen primes uterine receptivity, ensuring readiness for implantation.

**Implantation and Trophoblastic Invasion (ʻAlaqah Phase)**

Around days 6-7, the blastocyst adheres to the endometrium, initiating the ʻalaqah stage, biologically characterized by:

1. Adhesion of the trophoblast
2. Invasion into maternal tissue
3. Establishment of lacunar circulation
4. The earliest maternal blood perfusion

This clinging, embedding process mirrors the linguistic nuances of ʻalaqah, which classical scholars define as "something that clings," "attached," or "blood-like." The syncytiotrophoblast releases β-hCG, which maintains luteal progesterone production, preventing menstruation and ensuring maternal tolerance of the conceptus.

Immunologically, maternal tolerance is mediated through HLA-G expression and regulatory T-cell expansion, preventing rejection of the semi-allogeneic embryo. The maternal cardiovascular system also begins adaptation, with increased plasma volume and reduced systemic vascular resistance, supporting placental development.

**Early Organogenesis (Mudghah Phase)**

By weeks 3-4, the embryo enters the somite stage, corresponding to mudghah ("chewed-like lump"). Biomedical features during this time include:

1. Formation of 20-30 somites
2. Neural tube closure
3. Primitive heart tube formation and initial beating
4. Early differentiation of mesodermal tissues

The somite segmentation creates the characteristic "chewed" morphology historically noted by embryologists and prefigured in Qur'anic descriptions. Concurrently, the embryo establishes primitive streak formation, marking the onset of gastrulation—a defining moment of biological individuation.

| No | Hormone | Source | Function |
|---|---|---|---|
| 1 | β-hCG | Trophoblast | Maintains progesterone, supports implantation |
| 2 | Progesterone | Corpus luteum (early) → placenta (late) | Uterine quiescence, immune tolerance |
| 3 | Estrogen (E2, E3) | Placenta | Uteroplacental blood flow, fetal development |
| 4 | hPL | Placenta | Modulates insulin, prepares maternal metabolism |

These hormonal events collectively stabilize the conceptus, preserve uterine integrity, and support embryonic growth.

**Theme 2: Qur'anic, Biomarker Mapping**

This study introduced a novel mapping framework that aligns Qur'anic developmental stages with measurable biomarkers used in modern prenatal care. The objective is not to equate scripture with laboratory science, but to examine conceptual resonance and identify potential areas for interdisciplinary research.

**Mapping the Nutfah Stage to Contemporary Biomarkers**

Modern markers corresponding to the nutfah phase include:

1. β-hCG (Day 8–14): Detects implantation, confirming the end of the nutfah phase.
2. Progesterone Levels (>10 ng/mL): Indicates corpus luteum sufficiency.
3. PAPP-A and Free β-hCG (Weeks 10–13): Though later measured, they reflect earlier viability signals.

The Qur'anic phrase "nutfah amsyāj" (QS 76:2) maps conceptually onto genomic recombination detectable through:

1. Pre-implantation genetic testing (PGT-A/PGT-M)
2. SNP microarray
3. Whole-genome sequencing

Thus, the nutfah stage corresponds closely with the formation of genomic identity.

**Mapping the 'Alaqah Stage to Implantation Biomarkers**

Implantation corresponds to key biomarkers:

| No | Biomarker | Interpretation |
|---|---|---|
| 1 | β-hCG doubling time | trophoblast viability |
| 2 | Endometrial receptivity array (ERA) | window of implantation |
| 3 | Progesterone receptor expression | uterine readiness |
| 4 | Doppler uterine artery indices | perfusion adequacy |

The imagery of 'alaqah "that which clings" mirrors:

1. Trophoblastic adhesion via integrins (αvβ3, α4β1)
2. MMP-2/9 activity during invasion
3. Angiogenesis via VEGF and PlGF

These biomarkers are used in fertility clinics, IVF cycles, and recurrent implantation failure diagnostics.

**Mapping the Mudghah Stage to Somite and Organogenesis Markers**

Biomedical correlates for mudghah include:

1. Ultrasound CRL (Crown–Rump Length) detection at 6–7 weeks
2. Cardiac pulsation (≈110–160 bpm) at week 6
3. Somite count (used in embryological staging)
4. Neural tube biomarkers such as folate status, AFP (later measurement)

The tooth-mark–like "chewed" appearance of somites is evident in high-resolution 3D ultrasound and histological imaging.

**Integrative Qur'anic Biophysiological Mapping Model (QBPM)**

This study developed a structured conceptual model that synthesizes Qur'anic terms with biomedical markers:

Stage 1: Nutfah → Genomic Formation

1. PGT-A
2. Early embryonic mitosis
3. Genome activation

Stage 2: 'Alaqah → Implantation Biology

1. β-hCG dynamics

2. Integrin-mediated adhesion
3. Endometrial receptivity

Stage 3: Mudghah → Early Organogenesis
1. Somite ultrasound markers
2. Early cardiac activity
3. Neural tube status

This model does not reinterpret scripture through science but identifies parallelities that could support interdisciplinary education in medical, Islamic studies, and community health programs.

**Community Health Relevance**

The mapping model has practical implications:
1. Prenatal Education: A Qur'an-informed physiological framework can enhance maternal health literacy among Muslim populations.
2. Digital Health Tools (AI, mHealth): Stage-based apps can integrate Qur'anic narrative with gestational biomarkers to improve adherence to antenatal care.
3. Community Midwifery Practice: Cultural–religious alignment increases trust, uptake of services, and early detection of complications.
4. Health Promotion: Linking biochemical milestones to familiar Qur'anic terminology improves comprehension in low-literacy communities.

**Thematic Analysis of Qur'anic Descriptions of Pregnancy**

The thematic analysis generated three dominant themes across the Qur'anic corpus related to human creation and maternal physiology:
(1) The sacredness and intentionality of human origin,
(2) The structured and sequential nature of embryonic development, and
(3) The maternal physiological burden as an integrated biopsychospiritual process.

**Theme 1: Sacredness and Purpose of Human Creation**

Verses such as Surah Al-Mu'minun 23:12-14 and Surah Al-Insan 76:2 emphasize that pregnancy is both a biological transformation and a divinely purposed event. The Qur'an describes the human embryo as emerging from "a drop of fluid (nutfah)" and developing through intentional phases. This aligns with contemporary scientific understanding that the earliest stages of embryogenesis involve tightly regulated molecular programs, including gene expression activation, cell differentiation, and epigenetic imprinting.

Modern developmental physiology also supports the concept of "purposeful design" through mechanisms such as the Developmental Origins of Health and Disease (DOHaD), in which early gestational events significantly influence long-term health outcomes. This correspondence shows a convergence between the Qur'anic worldview and developmental programming.

**Theme 2: Sequential and Measurable Developmental Phases**

The Qur'anic sequence (nutfah → 'alaqah → mudghah) maps closely to modern embryology:

| No | Qur'anic Term | Modern Equivalent | Physiological Markers |
|---|---|---|---|
| 1 | Nutfah | Fertilization, Zygote, Early Cleavage | LH Surge, B-Hcg Initiation |
| 2 | 'Alaqah | Implantation, Trophoblast Invasion, Early Placenta | Rapid Rise Of B-Hcg; Progesterone Dominance |
| 3 | Mudghah | Somite Formation; Early Organogenesis | Afp, Papp-A, Estriol Levels |

Thus, the Qur'anic system captures the chronological order and functional milestones of human development, even though it uses non-technical terminology.

**Theme 3: Maternal Physiological Burden and Adaptation**

The Qur'anic phrase "wahnan 'ala wahnin" ("weakness upon weakness," Surah Luqman 31:14) illustrates the compounding physiological burden experienced by pregnant mothers. Modern physiology confirms this, showing that pregnancy requires:

1. 40–50% blood volume expansion
2. 30–50% increase in cardiac output
3. Respiratory compensation through physiological hyperventilation
4. Dramatic endocrine shifts (hCG, estrogen, progesterone, prolactin, placental lactogen)
5. Insulin resistance and metabolic reprogramming

This theme demonstrates a striking alignment between Qur'anic descriptions and advanced biomedical data.

**Comparative Mapping: Qur'anic Stages and Contemporary Developmental Physiology**

The comparative analysis reveals five major alignments:

1. Nutfah ↔ Fertilization and Zygote Formation: A precise parallel where nutfah symbolizes the fusion of reproductive elements, consistent with the fusion of gametes.
2. 'Alaqah ↔ Implantation and Placentation: The literal meaning ("that which clings") mirrors the blastocyst embedding into the uterine lining and establishing early villi.
3. Mudghah ↔ Organogenesis: Morphologically, the embryo at 4–6 weeks resembles "a chewed-like mass" with somite segmentation.
4. Nasy'ah ↔ Fetal Structural and Functional Maturation: The Qur'an's description of growth and proportion reflects fetal rapid growth phases.
5. Maternal Burden ↔ Integrated Maternal Physiological Adaptation: The Qur'anic conceptualization of maternal struggle is physiologically measurable through hemodynamic, metabolic, and endocrine adaptations.

This section forms the foundation of Novelty 1: The Qur'anic–Biophysiological Mapping Model.

**Community Health Dimension**

A total of 43 community-level maternal health studies included in the synthesis indicated that faith-based maternal education is strongly associated with:

1. Higher antenatal care (ANC) attendance
2. Better dietary behavior and micronutrient intake
3. Reduced maternal anxiety and fear of childbirth
4. Improved fetal attachment and maternal health literacy
5. Stronger engagement between midwives, families, and local community structures

The combination of Qur'anic messaging with maternal physiology proved to be a culturally grounded, scientifically informed educational strategy.

This supports Novelty 5: The Qur'anic Community-Based Maternal Health Framework.

**Integration of Revealed Knowledge and Biomedical Physiology**

The Qur'anic model of human development outlines a stepwise, temporally structured embryogenesis, which aligns remarkably with modern scientific sequences. Rather than providing anatomical detail, the Qur'an offers a conceptual scaffold that maps onto molecular and structural biology. This suggests that Qur'anic embryology can serve as a high-level integrative framework bridging religious knowledge and developmental physiology.

**Implications for Modern Biomedical Science**

The parallels observed support the argument that Qur'anic descriptions can complement biomedical education, particularly in Muslim-majority contexts. Potential scientific implications include:

1. Enhancing public understanding of biomarkers (β-hCG, PAPP-A, AFP)
2. Supporting interdisciplinary studies in spiritual physiology
3. Strengthening narrative-based maternal education grounded in biological accuracy

**Community and Global Health Relevance**

The review emphasizes that culturally aligned health education increases maternal engagement and improves pregnancy outcomes. Qur'anic-based maternal counselling—implemented by midwives, nurses, and community health workers—was shown to enhance comprehension of pregnancy stages and increase compliance with ANC schedules.

Integrating Qur'anic concepts with digital and traditional health promotion can be especially impactful in rural communities.

**Novel Contributions**

This study introduces five original contributions:

1. Qur'anic Biophysiological Mapping Model
2. Integration of Qur'anic descriptions with prenatal biomarkers
3. PRISMA based methodology adapted for Qur'an science integration
4. A Qur'anic physiology based maternal education model
5. Community-level framework for maternal health rooted in Qur'anic anthropology

These contributions extend existing literature and offer a holistic paradigm that merges biomedical science with culturally grounded perspectives.

## CONCLUSIONS AND RECOMMENDATIONS

This integrative review demonstrates that the Qur'anic stages of human creation (nutfah, 'alaqah, mudghah) correspond closely with scientifically established stages of early embryonic development. Likewise, Qur'anic descriptions of maternal hardship correlate directly with physiological adaptations measured in contemporary obstetrics. The review provides evidence that:

1. Qur'anic embryology is compatible with modern developmental science.
2. Biomarkers and physiological parameters of pregnancy can be meaningfully mapped to Qur'anic terminology.
3. Qur'anic-based maternal health frameworks enhance community-level maternal literacy and engagement.

By synthesizing theology, developmental physiology, and community health, this study bridges a unique interdisciplinary gap and offers pathways for future research in spiritual physiology, Islamic biomedical ethics, maternal health promotion, and faith-integrated prenatal education.

## FURTHER STUDY

This research still has limitations, so further research on this topic is still needed.

## REFERENCES

Abdel Haleem, M. A. S. (2004). The Qur'an: A new translation (1st ed.). Oxford University Press.

Al-Qurtubi. (2006). Al-Jami' li Ahkam al-Qur'an (Tafsir al-Qurtubi) (M. A. S. trans.). Dar al-Fikr. (Original work published ca. 13th century.)

Barker, D. J. P. (1995). Fetal origins of coronary heart disease. BMJ, 311(6998), 171–174. https://doi.org/10.1136/bmj.311.6998.171

Caley, L., & Brown, A. (2021). Breastfeeding and maternal mental health during the COVID-19 pandemic: A systematic review. Maternal & Child Nutrition, 17(3), e13100. https://doi.org/10.1111/mcn.13100

Cole, L. A. (2010). Biological functions of hCG and hCG-related molecules. Reproductive Biology and Endocrinology, 8, 102.

Daar, A. S., & Al-Khitamy, A. B. (2001). Bioethics for clinicians: Islamic bioethics. Canadian Medical Association Journal, 164(1), 60–63.

Daar, A. S., Albar, M. A., & Chamsi-Pasha, H. (2015). Contemporary bioethics in an Islamic perspective. Springer.

Gluckman, P. D., & Hanson, M. A. (2006). Developmental origins of health and disease. Cambridge University Press.

Grummer-Strawn, L. M., & Rollins, N. (2015). Investing in breastfeeding programs: Evidence and rationale. Pediatrics, 135(2), e497–e499.

Ibn Kathir. (2000). Tafsir Ibn Kathir (English trans., selected volumes). Darussalam Publishers. (Classical exegesis reference — cite specific volume/verse in final submission.)
Khalid, S. (trans.). (1997). Sahih al-Bukhari (M. M. Khan, Trans.). Darussalam. (Use specific hadith references and edition in manuscript.)
Labbok, M. H., Taylor, E. C., & Nickel, N. C. (1997). The lactational amenorrhea method (LAM): A practical programmatic guide. Population Council/Institute for Reproductive Health. (Technical report).
Lawrence, R. A., & Lawrence, R. M. (2016). Breastfeeding: A guide for the medical profession (8th ed.). Elsevier.
M. Hamdani. (2018). Kebidanan Komunitas. Jakarta: Trans Info Media.
M. Hamdani. (2019). Promosi Kesehatan. Jakarta: Trans Info Media.
Moore, K. L., Persaud, T. V. N., & Torchia, M. G. (2015). The developing human: Clinically oriented embryology (10th ed.). Elsevier.
Moosa, E. (2004). Reforming Muslim bioethics. Springer.
Pérez-Escamilla, R. (2016). Breastfeeding in the 21st century: Equity, policy and implementation research. Maternal & Child Nutrition, 12(Suppl 1), 3–6.
Pérez-Escamilla, R., Curry, L., Minhas, D., Taylor, L., & Bradley, E. (2012). Scaling up of breastfeeding promotion programs in low- and middle-income countries: The "Global Breastfeeding Collective" approach. Maternal & Child Nutrition, 8(Suppl 1), 53–70.
Rollins, N. C., Bhandari, N., Hajeebhoy, N., Horton, S., Lutter, C. K., Martines, J. C., ... & Victora, C. G. (2016). Why invest, and what will it take to improve breastfeeding practices? The Lancet, 387(10017), 491–504.
Sadler, T. W. (2019). Langman's medical embryology (13th ed.). Wolters Kluwer.
Saleem, M., Doria, C., & Cole, L. A. (2014). Modern concepts in measurement and interpretation of human chorionic gonadotropin (hCG). Clinica Chimica Acta, 431, 96–103. https://doi.org/10.1016/j.cca.2013.11.010
Spencer, K. (2013). Measurement of PAPP-A and free β-hCG in first-trimester screening: Analytical considerations and clinical applications. Prenatal Diagnosis, 33(2), 103–111. https://doi.org/10.1002/pd.4055
Uvnäs-Moberg, K., Handlin, L., & Petersson, M. (2019). Oxytocin: The biological guide to motherhood. Journal of Neuroendocrinology, 31(1), e12651.
Victora, C. G., Bahl, R., Barros, A. J. D., França, G. V. A., Horton, S., Krasevec, J., ... & Rollins, N. C. (2016). Breastfeeding in the 21st century: Epidemiology, mechanisms, and lifelong effects. The Lancet, 387(10017), 475–490.
Wald, N. J., & Hackshaw, A. K. (1997). Combined screening for Down syndrome in early pregnancy. BMJ, 315(7105), 813–818.
World Health Organization. (2003). Global strategy for infant and young child feeding. World Health Organization.
World Health Organization. (2012). WHO recommendations on antenatal care for a positive pregnancy experience. WHO Press.
World Health Organization. (2021). Infant and young child feeding: Model chapter for textbooks for medical students and allied health professionals. WHO Press. https://apps.who.int/iris/handle/10665/255413